

Corrigenda: *Stygobromus bakeri*, a new species of groundwater amphipod (Amphipoda, Crangonyctidae) associated with the Trinity and Edwards aquifers of central Texas, USA. Subterranean Biology 38: 19–45. <https://doi:10.3897/subtbiol.38.61787>

Randy Gibson¹, Benjamin T. Hutchins², Jean K. Krejca³,
Peter H. Diaz⁴, Peter S. Sprouse³

1 San Marcos Aquatic Resources Center, United States Fish and Wildlife Service, 500 East McCarty Lane, San Marcos, Texas, 78666, USA **2** Edwards Aquifer Research and Data Center, Texas State University, 220 East Sessom Drive, San Marcos, Texas 78666, USA **3** Zara Environmental LLC., 1707 West FM 1626, Manchaca, TX, 78652, USA **4** Texas Fish and Wildlife Conservation Office, United States Fish and Wildlife Service, 500 East McCarty Lane, San Marcos, Texas, 78666, USA

Corresponding author: Randy Gibson (randy_gibson@fws.gov)

Academic editor: O.T. Moldovan | Received 12 May 2021 | Accepted 14 May 2021 | Published 21 May 2021

<http://zoobank.org/11BDC2F7-406D-408D-A12A-BB01C836C4D6>

Citation: Gibson R, Hutchins BT, Krejca JK, Diaz PH, Sprouse PS (2021) Corrigenda: *Stygobromus bakeri*, a new species of groundwater amphipod (Amphipoda, Crangonyctidae) associated with the Trinity and Edwards aquifers of central Texas, USA. Subterranean Biology 38: 19–45. <https://doi:10.3897/subtbiol.38.61787>. Subterranean Biology 38: 113–115. <https://doi.org/10.3897/subtbiol.38.68597>

Keywords

subterranean, karst, stygobiont, stygobite, troglomorphy, crenobiont, interstitial crustacean, spring, cave, sympatric

Introduction

It has come to our attention that many of the records reported in Table 1 were shifted or missing and therefore incorrect.

The CORRECT Table is as follows:

Table 1. Groundwater fauna collected at each *Stygobromus bakeri* sp. nov. site. Stygobionts (S) are hypogean and colonize various groundwater habitats. Crenobionts (C) are epigean and preferentially colonize springs.

| TAXON | STYGOBIONT/ CRENOBIOT | JOHN KNOX SPRING | JACOB'S WELL | SALADO SPRINGS | MORMON SPRING #3 |
|---|--------------------------|---------------------|--------------|-------------------|---------------------|
| Turbellarians | | | | | |
| Order Tricladida | | | | | |
| Family Kenkiidae | | | | | |
| <i>Sphallopiana mohri</i> Hyman, 1938 | S | X | | X | |
| Annelids | | | | | |
| Order Lumbriculata | | | | | |
| Family Lumbriculidae | | | | | |
| <i>Eremidrilus</i> sp. 1 Fend & Rodriguez, 2003 | S | X | | X | |
| <i>Eremidrilus</i> sp. 2 | S | | | X | |
| Molluscs | | | | | |
| Order Gastropoda | | | | | |
| Family Cochliopidae | | | | | |
| <i>Phreatoceras taylori</i> (Hershler & Longley, 1986a) | S | | | X | |
| <i>Phreatodrobia conica</i> Hershler & Longley, 1986b | S | | | X | |
| <i>Phreatodrobia micra</i> (Pilsbry & Ferriss, 1906) | S | | | X | |
| <i>Phreatodrobia nugax</i> (Pilsbry & Ferriss, 1906) | S | X | | X | |
| <i>Stygopyrgia bartonensis</i> Hershler & Longley, 1986b | S | | | | X |
| Family Hydrobiidae | | | | | |
| <i>Marstonia comalensis</i> (Pilsbry & Ferriss, 1906) | C | X | X | X | |
| Family Plueroceridae | | | | | |
| <i>Elimia comalensis</i> (Pilsbry, 1890) | C | | X | X | |
| Crustaceans | | | | | |
| Order Amphipoda | | | | | |
| Family Bogidiellidae | | | | | |
| <i>Parabogidiella americana</i> Holsinger, 1980 | S | | | X | |
| Family Crangonyctidae | | | | | |
| <i>Stygobromus bifurcatus</i> (Holsinger, 1967) | S | | X | X | X |
| <i>Stygobromus russelli</i> (Holsinger, 1967) | S | X | X | X | X |
| <i>Stygobromus (flagellatus group)</i> sp. nov. Cope, 1872 | S | | X | | |
| Order Bathynellacea | | | | | |
| Family Parabathynellidae | | | | | |
| <i>Texanobathynella bowmani</i> Delamare Deboutteville, Coineau & Serban, 1975 | S | | | X | |
| Order Isopoda | | | | | |
| Family Asellidae | | | | | |
| <i>Caecidotea bilineata</i> Lewis & Bowman, 1996 | S | | | X | |
| <i>Caecidotea reddelli</i> (Steeves, 1968) | S | | | X | X |
| <i>Lirceolus bisetus</i> (Steeves, 1968) | | | | X | |
| <i>Lirceolus hardeni</i> (Lewis & Bowman, 1996) | S | X | X | X | |
| <i>Lirceolus pilus</i> (Steeves, 1968) | S | | X | X | |
| Order Podocopida | | | | | |
| Family Candonidae | | | | | |
| <i>Comalcandona tressleri</i> Külköylüoğlu & Gibson, 2018 | S | X | | | |
| <i>Schornikovdona bellensis</i> Külköylüoğlu, Yavuzatmaca, Akdemir, Diaz & Gibson, 2017 | S | | | X | |
| <i>Pseudocandona albicans</i> (Brady, 1864) | C | X | | | |
| <i>Pseudocandona cf. semicognita</i> (Schäfer, 1934) | C | X | | | |
| <i>Physocypris cf. globula</i> Furtos, 1933 | C | X | | | |
| <i>Darwinulopsis stevensoni</i> (Brady & Robertson, 1870) | C | X | | | |
| <i>Cypridopsis</i> sp. (Brady, 1867) | C | X | | | |
| <i>Physocypris denticulata</i> (Daday, 1905) | C | | | | X |
| <i>Chlamydotheca texasiensis</i> (Baird, 1862) | C | | | | X |
| <i>Cypridopsis cf. helvetica</i> Kaufmann, 1900 | C | X | | | |

| Taxon | Stygobiont/ Crenobiont | John Knox Spring | Jacob's Well | Salado Springs | Mormon Spring #3 |
|---|---------------------------|---------------------|--------------|-------------------|---------------------|
| Arachnids | | | | | |
| Order Trobidiformes | | | | | |
| Family Arrenuridae | | | | | |
| <i>Arrenurus</i> spp. Dugès, 1834 | S | X | | X | |
| Insects | | | | | |
| Order Coleoptera | | | | | |
| Family Dytiscidae | | | | | |
| <i>Sanilippodytes</i> sp. Franciscolo, 1979 | C | X | | X | |
| <i>Uvarus texanus</i> (Sharp, 1882) | C | | | X | |
| Family Elmidae | | | | | |
| <i>Microcloepus pusillus</i> (LeConte, 1852) | C | X | X | X | X |
| Salamanders | | | | | |
| Order Urodela | | | | | |
| Family Plethodontidae | | | | | |
| <i>Eurycea chisholmensis</i> Chippindale, Price, Wiens & Hillis, 2000 | C | | | X | |
| <i>Eurycea pterophila</i> Burger, Smith & Potter, 1950 | C | X | X | | |