

# A new species of *Troglobius* (Collembola, Paronellidae, Cyphoderinae) from a Brazilian iron cave

Douglas Zeppelini<sup>1,4,†</sup>, Diego Dias da Silva<sup>2,‡</sup>, José G. Palacios-Vargas<sup>3,§</sup>

**1** Laboratório de Sistemática de Collembola e Conservação. Depto. Biologia, Centro de Ciências Biológicas e Sociais Aplicadas. Universidade Estadual da Paraíba campus V, 58070-450, João Pessoa, PB, Brazil **2** Programa de Pós Graduação em Sistemática e Evolução, Centro de Biociências, Universidade Federal do Rio Grande do Norte. 59072-970, BR 101, Campus Universitário, s/n, Lagoa Nova, Natal - RN - Brazil **3** Laboratorio de Ecología y Sistemática de Microartrópodos, Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Universidad Nacional Autónoma de México, 04510 México **4** Associação Guajiru-Ciência-Educação-Meio Ambiente. Scientific Board, 58310-000 João Pessoa, PB, Brazil

Corresponding author: Douglas Zeppelini (zeppelini@daad-alumni.de)

Academic editor: O. Moldovan | Received 22 February 2014 | Accepted 28 April 2014 | Published 20 June 2014

<http://zoobank.org/0617F669-19D5-4A90-B37B-0587C3D9239E>

**Citation:** Zeppelini D, da Silva DD, Palacios-Vargas JG (2014) A new species of *Troglobius* (Collembola, Paronellidae, Cyphoderinae) from a Brazilian iron cave. Subterranean Biology 14: 1–13. doi: 10.3897/subtbiol.14.7355

## Abstract

The genus *Troglobius* (Collembola: Paronellidae: Cyphoderinae) is known only from the Southern Hemisphere, the two species found so far were described from caves in Madagascar and Brazil. In this paper we describe a new Brazilian species and complement the descriptions of the previously known species. The new species is a troglomorph and was collected in a single cave, in iron formations in the state of Minas Gerais. Brazil. The original diagnosis of the genus is supplemented with new characters based on observations of the new species.

## Keywords

Cave Fauna, Iron Ore, Iron Caves, Taxonomy, Southern Hemisphere

## Introduction

The genus *Troglobius* Palacios-Vargas & Wilson, 1990 comprises two species, both from caves in Southern Hemisphere, the type species *T. coprophagus* from Madagascar and *T. brasiliensis* Palacios-Vargas & Zeppelini, 1995 from Brazil. The genus belongs

to the Collembola family Paronellidae Börner, 1913 (Entomobryomorpha) and has been assigned to the subfamily Paronellinae since its original description and in recent contributions (Soto-Adames et al. 2008).

The distribution of the genus in Brazil ranges from caves in the Amazon rain forest in Northern region, to caves in the subtropical Atlantic Forest domains in Southeastern region. Sampling in caves in iron formations are bringing new records to the distribution of the fauna in Brazilian iron caves, and important information on the generic characters of the genus. Unpublished material is still waiting for formal descriptions suggesting that the diversity of the genus in Brazil is likely to be much greater than the single described species.

The gap of information about cave fauna and cave Collembola from Africa make for the moment difficult to propose biogeographic hypothesis to explain the origin and distribution of the genus. Considering that the two new species (*T. coprophagus* and *T. brasiliensis*) are confined to caves (as it is known up to date) and isolated from one another with disjoint distribution, we can suppose that this is a relict genus with Gondwanian distribution. However, the answer will depend on the distribution and better knowledge of the African species.

This study presents a new species of *Troglobius* from iron-caves, which is described and compared with the two already described species. Furthermore, complementary morphological data is given to the description of *T. coprophagus* and *T. brasiliensis*. A new diagnosis is proposed to the genus, the taxonomy was revised and comparative analysis of the morphology of the species, with the diagnosis of subfamilies in Paronellidae resulted in new taxonomic arrangement for the genus in the Cyphoderinae subfamily.

## Methodology

Specimens were collected with pitfall traps, fixed in ethanol 70% and slide mounted in Hoyer' solution. Two paratypes were dissected. Head and body chaetotaxic systems followed Soto-Adames et al. (2008) and Szeptycki (1979). Abbreviations used in the text: Ant I-IV, antennal segments I-IV; Th II-III, thoracic segments II-III; Abd I-VI, abdominal segments I-VI.

## Redefinition of the genus *Troglobius* Palacios-Vargas & Wilson, 1990

### Taxonomic position:

Family Paronellidae Börner, 1913

Subfamily Cyphoderinae Börner, 1913 (sensu Soto-Adames et al., 2008) comb. n.

**Genus *Troglobius* Palacios-Vargas & Wilson, 1990**

**Type species.** *Troglobius coprophagus* Palacios-Vargas & Wilson, 1990

**Diagnosis.** Habitus as Entomobryomorpha, eyeless, without pigment. Body with ciliated chaetae and hyaline ovoid scales. Mesothorax and metathorax not humped or bent. Ant. IV not subdivided. Labial chaetotaxy with chaetae: M1, M2, R, E and L1 present, L2 absent. Head bearing two pairs of trichobothria. Mucro elongate, four times as long, or more, than its greatest width, with many serrations along one or both dorsal lamellae. Dens without spines, anterior side covered with scales. A couple of large scale, about the same size as mucro, present on apex of anterior surface of dens. Posterior side of dens with, at least, three rows of chaetae and scales. Some (or most) scales, or thick chaetae, of internal row deeply fringed. In other rows only some distal chaetae modified. Unguis and unguiculus with winged teeth.

**Remarks.** The original description places the genus in the family Paronellidae Börner, 1913. Soto-Adames et al. (2008) define the subfamilies Paronellinae and Cyphoderinae as follows:

*Separation between Paronellinae and Cyphoderinae is based on the presence of fringed dental scales in the latter and their absence in the former. All cyphoderines are eyeless, whereas the majority of paronellines have eyes. In addition, all cyphoderines have scales, whereas many paronellines lack these scales.*

The morphology of the dental scales, the absence of eyes, the shape of mucro and the presence of large scales at the tip of dens, about the same size as mucro, indicate that the genus *Troglobius* belongs to the subfamily Cyphoderinae (with 128 species distributed among 13 genera of blind Collembola) instead of Paronellinae (with 384 species in 27 genera) as proposed by Bellinger et al. (2013).

In the new proposed diagnosis for the genus, we conclude that the labial chaeta "R" is present, normal shaped, and variable in size, while chaeta L2 is always absent. An additional lateroventral trichobothria, close to the basis of antennae is present in each side of head, this feature is also seen in *Cyphoderus*, another genus of Cyphoderinae. The genus *Troglobius* can be differentiated from other Cyphoderinae by the shape of the mucro, which has, at least one lamella serrated.


**Description of the new species**

***Troglobius ferroicus* sp. n.**

<http://zoobank.org/C04718C0-B0D7-42F8-9304-96D66BEDFA11>

Figs 1–13

**Type locality.** Brazil, Minas Gerais, Itabirito, Várzea do Lopes. Cave VL29/30, 20°20'05"S, 43°56'18"W. Aphotic zone on water film over the substrate and organic matter, 8 individuals collected.




**Figures 1–2.** 1 Habitus, *Troglobius ferroicus* sp. n. 2 Antennal sensilla of apex of Ant. II, same type of sensilla seen on apex of Ant. III. *T. ferroicus* sp. n.

*Holotype.* Slide mounted in Hoyer's solution. Deposited at “Coleção de Referência de Fauna de Solo da Paraíba – Universidade Estadual da Paraíba” (CRFS/UEPB), #2766. Várzea do Lopes, Itabirito, MG, Brazil. Cave VL29/30. 02-06/x/2011. Leg. Speleology team of Carste consultants.

*Paratypes.* One paratype deposited at Laboratorio de Ecología y Sistemática de Microartrópodos - UNAM (slide #2837 donated from CRFS/UEPB). Várzea do Lopes, Itabirito, MG, Brazil. Cave VL 29/30. Andrade Col. 03-20/XI/2007.

One paratype deposited at Museu Nacional – Universidade Federal do Rio de Janeiro - MN/UFRJ (slide #2765 donated from CRFS/UEPB) Várzea do Lopes, Itabirito, MG, Brazil. Cave VL 29/30. 03/IV/2012. Other two paratypes at CRFS/UEPB, under numbers #2838 and 3098, from the same locality as the holotype.




**Figure 3.** Cephalic chaetotaxy A (*T. ferroicus* sp. n.), B (*T. brasiliensis*), C (*T. coprophagus*).

**Description.** Habitus of entomobryomorphs (Fig. 1). Length (n = 4) 2.3 mm in average (1.9–2.7 mm). Color pearly white. Body with smooth and ciliated chaetae of different sizes, oval scales hyaline, trichobothria long and ciliated.

Ratio head:antenna 1:2.5, ratio antenna:body length 1:1.9. Ant. I–II with scales. Ratio of Ant. I–IV as 1: 2.6; 2.5; 3.7. Apex of Ant. II with three candle-shaped sensilla, two close to each other and a third laterally displaced (Fig. 2). Apical sensory organ of Ant. III with two candle-shaped sensilla, apical guard chaetae thicker than others. Ant. IV undivided, with rows of chaetae and sensilla or finely ciliated microchaetae, without apical bulb.


Eyes and pigment absent, head with two pairs of trichobothria (Fig. 3A), chaetae Ps5 and Pa5 ciliated. Pre-labral and labral chaetae formula 4/5, 5,4; all chaetae



**Figures 4–8.** **4** Labial triangle, *T. ferroicus* sp. n. **5** Metatrochanteral organ, *T. ferroicus* sp. n. **6** Unguis I, *T. ferroicus* sp. n. **7** Unguis II, *T. ferroicus* sp. n. **8** Unguis III, *T. ferroicus* sp. n.

smooth. Labial triangle with anterior row (A1–5) smooth. All labial triangle chaetae smooth, labial seta R well developed, L2 absent (Fig. 4).

Legs with ciliated setae. Coxa of leg I with one chaeta. Metatrochanteral organ with 21 chaetae forming a “V” and 13 additional short chaetae (Fig. 5). Tenent hairs acuminate, shorter than ventral lamella of unguis (Fig. 6–8); ratio unguis: tenent hair = 1: 0.65. All unguis with two inner basal winged teeth of different sizes; one median inner tooth and one very small tooth at the base of the outer lamella. All unguiculi with well-developed outer lamella and normal sized inner lamella (Figs 6–8).




**Figure 9.** Body chaetotaxy *T. ferroicus* sp. n.

Abd II with two trichobothria, Abd III-IV with three trichobothria each; Abd. IV with 4+4 lenticular organs on posterior margin (Fig. 9). Posterior surface of ventral tube with 2 long macrochaetae, scales on proximal end, 10 proximal pairs of ciliated chaetae and 11 distal pairs of small smooth chaetae; anterior face with only 2 pairs of ciliated macrochaeta (Fig. 10).

Furcula with scales and ciliated chaetae; mucro, dens and manubrium ratio 1.0:3.8:4.2. The distal end of dens with a pair of large scales, about the same size of mucro (Fig. 11). Dens without spines, normal scales on anterior side, internal row of posterior face with 21 fringed scales (Fig. 12A), median row with one fringed scale (Fig. 13A). Other rows with finely ciliated chaetae in irregular pattern. Mucro elongated, bearing 10–15 teeth on outer lamella, inner lamella smooth with one tooth at the tip.

**Etymology.** The name of the new species *T. ferroicus* sp. n. comes from Latin *ferum* (iron) and Greek *oikos* (house). This makes allusion to the iron caves where the new species was found.



**Figures 10–11.** 10 Ventral tube chaetotaxy, lateral view, *T. ferroicus* sp. n. 11 Distal end of dens and mucro, *T. ferroicus* sp. n.


**Distribution.** Good's biogeographic zone 27, northeastern and central Brazil (Good 1974, Culik and Zeppelini 2003). The climate is defined by dry winters and wet summers, with average temperatures of 18 °C during winter and 22 °C in summer.

**Habitat.** *T. ferroicus* sp. n. is found in small (less than 100 m deep) iron-caves, dwelling on water film over the rocks and organic debris in the aphotic zone. The species was found in a single cave, and it is likely to be a troglobite which lives in the in the Mesovoid Shallow Substratum (MSS) (Juberthie et al. 1980) and the adjacent cave.

**Conservation status.** This cave is placed in a mining area with iron ores, which represents a putative threat to the species if the cave and surrounding environments, including the MSS, are not preserved. The species was included in the recent evaluation of the Brazilian Fauna Red List, and must be published in the next issue of the Brazilian Red List as Critically endangered (CR).

**Remarks.** *T. ferroicus* sp. n. is the largest species in the genus, adults are 2.3 mm long in average (2–2.8 mm), while other species are about 1.5 mm long (Table 1). It has the longest antennae and ratio of head:antennae (Table 1). Ungual internal unpaired teeth are lacking in *T. coprophagus*, there are two teeth in *T. brasiliensis* and a single tooth in *T. ferroicus* sp. n. Mucronal serration is different in *T. brasiliensis*, where both edges are serrated, in *T. coprophagus* and *T. ferroicus* sp. n. the inner edge is smooth. One important diagnostic feature to differentiate the species is the relative size of mucro and dens, the ratio mucro/dens in *T. ferroicus* sp. n. is 1:4.0, in *T. brasiliensis* is 1:3.0 and *T. coprophagus* 1:2.5. Number and shape of the dental scales can, also, easily differentiate the species (Figs 12–13). Chaeta m2 on Abd. I is lacking in *T. ferroicus* sp. n., and present in the other two species, and chaeta T1 on Abd. IV is present in *T. ferroicus* sp. n., but absent in the other two species.





**Figures 12–13.** 12 Dental chaetae A (*T. ferroicus* sp. n.), B (*T. brasiliensis*), C (*T. coprophagus*) 13 Dens and mucro A (*T. ferroicus* sp. n.), B (*T. brasiliensis*), C (*T. coprophagus*).

**Table 1.** Comparative measurements and ratios of the three species of the genus *Troglobius*.

	Size	Ratio head/Ant	Ratio body/Ant	Ratio Ant segments	Ungual teeth	Ratio mucro/ dens;man
<i>T. brasiliensis</i>	1.5	1:1.8	1:1.7	1:2.5; 2.5; 3.0	2	1:3.5;4.2
<i>T. coprophagus</i>	1.4	1:1.7	1:2.3	1:1.2; 5.6; 13.1	0	1:3.4;5.7
<i>T. ferroicus</i> sp.n.	2.3	1:2.5	1:1.9	1:2.6; 2.5; 3.7	1	1:3.8;4.2

Finally, *T. coprophagus* was found in La grotte d'Andrafiabe (25°C and 100% of humidity) in Ankarana, which is a small limestone massif in the extreme North of Madagascar. *T. brasiliensis* was found in Cave Limoeiro at Pará, North Brazil and Gruta dos Paiva at Sao Paulo, Southeastern Brazil, both in limestone massifs. The new species *T. ferroicus* sp. n. was found in one iron cave in the state of Minas Gerais, with very different climate and environmental conditions.


## Complements to the description of previous species

### *Troglobius brasiliensis* Palacios-Vargas & Zeppelini, 1995

**Remarks.** Head chaetotaxy as Fig. 3B. Apex of Ant. II and Ant. III with small candle shaped sensillae. Dens anterior side with hyaline scales; posterior side with three rows of modified thick chaetae (Fig. 12B), internal row with 21 fringed chaetae, the distal chaeta of median row is also heavily fringed, external row with about 20 barbulate chaetae (Fig. 13B). A large chaeta, about as long as mucro, is present at the distal tip of the dens. Mucro with 12–14 teeth on inner lamella and 14 in outer lamella. Body chaetotaxy as in Fig. 14.

**Material examined.** Paratype, Brasil, Sao Paulo, Iporanga. Cave Gruta dos Paiva. col. Trajano, E. 20/II/1989.


It is important to mention that the species *T. brasiliensis* was described from specimens collected in caves in Northern Brazil. The type locality of this species is Cave Limoeiro (7-X-1988, Brasil, Pará, Medicilandia, Caverna Limoeiro. Col. E. Trajano), a small municipality in Para state in the Amazonic Region. A single specimen, assigned as paratype, comes from a different locality in Southeastern Brazil, far away from the type locality. This is the specimen studied here. Some differences were observed in specimens from both localities, as mentioned in the original description (Palacios-Vargas and Zeppelini 1995) and as shown in Fig. 13B in this work and Fig. 9 in the original description. The specimens from the type locality have many fringed chaetae and two fringed distal scales on posterior side of dens. The specimen from Iporanga has only fringed chaetae. This suggests that the Holotype and Paratype may represent different species, therefore, a detailed revision of the holotype of *T. brasiliensis* is needed.



*Troglobius coprophagus* Palacios-Vargas & Wilson, 1990

One of the species author, Wilson, was not able to maintain specimens alive once they were taken outside the cave, which indicates that they are very sensitive to temperature changes (Palacios-Vargas and Wilson 1990).

**Material examined.** Two paratypes, Madagascar, Grote d'Andrafiabe. 08/IX/1986.



**Figure 15.** Body chaetotaxy *T. coprophagus*.

## Acknowledgments

The senior author was granted by CNPq #301803/2012-9. The specimens were collected by the speleology team of Carste consultants. Roniere A. Brito (Universidade Estadual da Paraíba) helped in slide mountings. Blanca Mejía Recamier (Universidad Nacional Autónoma de México) dissected the specimens. Felipe Soto-Adames revised the English version of the manuscript.

## References

- Bellinger P, Christiansen K, Greenslade P, Mitra SK, Janssens F (2013) Checklist of Collembola: Key to the genera of Paronellinae. <http://www.collembola.org/key/paronellinae.htm>
- Börner VC (1913) Die Familien der Collembolen. Sonderabdruck aus dem Zoologischen Anzeiger 61(7): 315–322.
- Culik MP, Zeppelini D (2003) Diversity and distribution of Collembola: Arthropoda: Hexapoda of Brazil. Biodiversity and Conservation 12: 1119–1143. doi: 10.1023/A:1023069912619
- Good R (1974) The geography of flowering plants 4<sup>th</sup> ed. Longman Group, London, 574 pp.

- Juberthie C, Delay B, Bouillon M (1980) Sur l'existence du milieu souterrain superficiel en zone non calcaire. Comptes Rendus de l'Académie des Sciences de la France D, 290: 49–52.
- Palacios-Vargas JG, Wilson JM (1990) *Troglobius coprophagus*, a new genus and species of cave Collembola from Madagascar, with notes on its ecology. International Journal of Speleology 19: 67–73. doi: 10.5038/1827-806X.19.1.6
- Palacios-Vargas JG, Zeppelini D (1995) A new species of *Troglobius* (Collembola, Paronellidae) from Brazil. International Journal of Speleology 23(3–4): 173–177.
- Soto-Adames F, Barra JA, Christiansen K, Jordana R (2008) Suprageneric Classification of Collembola Entomobryomorpha. Annals of the Entomological Society of America 101(3): 501–513. doi: 10.1603/0013-8746(2008)101[501:SCOCE]2.0.CO;2
- Szeptycki A (1979) Chaetotaxy of the Entomobryidae and its phylogenetical significance. Morphosystematic studies of Collembola IV. Polska Akademia Nauk, Zakład Zoologii Systematycznej Doswiadczałnej, Krakow-Poland, 219 pp.